

BRAIDED LINE AND COUNTING FIXED POINTS OF $GL(d, \mathbb{F}_q)$

P.J. CAMERON + S. MAJID

ABSTRACT. We interpret a recent formula for counting orbits of $GL(d, \mathbb{F}_q)$ in terms of counting fixed points as addition in the affine braided line. The theory of such braided groups (or Hopf algebras in braided categories) allows us to obtain the inverse relationship, which turns out to be the same formula but with q and q^{-1} interchanged (a perfect duality between counting orbits and counting fixed points). In particular, the probability that an element of $GL(d, \mathbb{F}_q)$ has no fixed points is found to be the truncated q -exponential

$${}_d e_q^{-\frac{1}{q-1}} = \sum_{i=0}^d \frac{(-1)^i}{(q^i - 1) \cdots (q - 1)}.$$

1. INTRODUCTION

Let $\Sigma = \{1, \dots, d\}$. It is known [1] that one may write the number F_j of orbits of $G \subseteq S_d$ acting on the set Σ_j of distinct j -tuples in terms of the probabilities p_i that an element of G has i fixed points. Writing generating functions

$$F = \sum_{j=0}^d F_j \frac{x^j}{j!}, \quad p(x) = \sum_{i=0}^d p_i x^i,$$

the two are related by

$$F(x) = p(x+1).$$

For example, when $G = S_d$, it is evident that $F_j = 1$ for all j and hence

$$p_0 = {}_d e^{-1} \equiv \sum_{i=0}^d \frac{(-1)^i}{i!}.$$

Recently, one of us[2] has considered a similar problem for $G \subseteq GL(d, \mathbb{F}_q)$ where $\mathbb{F}_q$ denotes the field of prime-power order q . Let

$$\Sigma_j = \{(a_1, \dots, a_j), \quad a_i \in \mathbb{F}_q^d, \quad a_i \text{ linearly indept.}\}, \quad L_j = \#\{\text{Orbits of } G \text{ in } \Sigma_j\},$$

and let P_i denote the probability that an element of G pointwise fixes exactly a subspace of $\mathbb{F}_q^d$ of dimension i . Then similarly by the orbit-counting lemma one has

(1)

$$L_j = \frac{1}{|G|} \sum_{g \in G} \#\{g - \text{fixed points in } \Sigma_j\} = \sum_{i=j}^d P_i (q^i - 1)(q^i - q) \cdots (q^i - q^{j-1}).$$

S.M. is a Royal Society University Research Fellow.

The generating function

$$(2) \quad L(x) = \sum_{j=0}^d \frac{L_j x^j}{(q^j - 1) \cdots (q - 1)}$$

is likewise written in terms of the P_i and q -binomial coefficients. The latter are well-known (the Gauss formula) to count the number of subspaces of a given dimension in an $\mathbb{F}_q$ -vector space. The q -factorial denominators in such expressions are understood as 1 for $j = 0$.

On the other hand, for applications, we would really like the inverse relationship P_i in terms of L_j . In this note we provide this inverse in a natural way. Thus:

Proposition 1.1. *With the notations above the probabilities P_i of an i -dimensional fixed subspace are given by*

$$P_j = \frac{q^{-\frac{j(j-1)}{2}}}{(q^j - 1) \cdots (q - 1)} \sum_{i=j}^d \frac{L_i (-1)^{i-j}}{(q^i - q^j) \cdots (q^{j+1} - q^j)}.$$

For example, it is evident that if $G = GL(d, \mathbb{F}_q)$ then $L_j = 1$ for all j . Hence we find:

Corollary 1.2. *For $G = GL(d, \mathbb{F}_q)$ the probability of no fixed points is*

$$P_0 = \sum_{i=0}^d \frac{(-1)^i}{(q^i - 1) \cdots (q - 1)} \equiv {}_d e_q^{-\frac{1}{(q-1)}}.$$

Our standard notation for the (truncated) q -exponential here is such that ${}_d e_q^x \rightarrow {}_d e^x$ as $q \rightarrow 1$, i.e. it is defined with the factorial of q -integers $[i]_q = \frac{q^i - 1}{q - 1}$. In our case $q \geq 2$ and these $(q - 1)$ factors cancel but the notation reminds us of a divergence if we were to try include the finite group case as $q = 1$.

There are of course many ways to invert a linear relationship. We mention that one may also, for example, use the method of Möbius inversion with respect to a suitable function, see [3, p. 127]. On the other hand we will obtain the above inversion in a way that is interesting because it is precisely analogous to the computation for the S_d case based on generating functions. The only difference is that that one should consider the generating function dummy variable x for the purposes of addition and subtraction as a braided one, i.e. the polynomials $k[x]$ as a Hopf algebra, but in a braided category where x has braid statistics q (the affine ‘braided line’). (The idea here is similar to that of a superspace but instead of a ‘fermionic’ factor -1 there is a factor q when x is interchanged with another independent x .) The systematic passage from permutation groups to linear groups over finite fields by means of switching on a braiding would appear to be an interesting direction suggested by the present work.

The required methods of Hopf algebras in braided categories or ‘braided groups’ have been introduced some years ago[4], precisely as the the basic structure underlying quantum groups and q -analysis, [5][6]. We recall what we need in a self-contained preliminary Section 2. Working with x as a braided variable we relate, in Section 3, the signed generating function $\bar{L}(x) = L(-x)$ and $P(x) = \sum_{i=0}^d P_i x^i$ by

$$(3) \quad \bar{L}(x) = P(y -_q x)|_{y=1},$$




FIGURE 1. Axioms of a braided group: (a) associativity of the product (similarly for Δ) and the braided homomorphism axiom, (b) the antipode axioms

where $-_q$ on the right is subtraction under the additive braided group structure of the braided line. We can then immediately invert the relationship as

$$(4) \quad P(x) = \bar{L}(y -_{q^{-1}} x)|_{y=1},$$

where $-_{q^{-1}}$ is subtraction under the opposite additive braided group structure (it is not commutative). This gives the above inversion formula, essentially without computation. It also exhibits a novel duality phenomenon whereby the coefficients of $P(x)$ are given in terms of those of $\bar{L}(x)$ by exactly the same formula as vice-versa, but with q and q^{-1} interchanged. This is discussed in Section 4.

2. PRELIMINARIES: THE AFFINE BRAIDED LINE

We briefly recall the definition of braided groups and the braided line. Once these are understood, the computation itself in the next section will be immediate.

The formal definition of a braided category is in [7]. It consists of a category C , a functor $\otimes : C \times C \rightarrow C$ and natural transformations $\Psi : \otimes \rightarrow \otimes^{\text{op}}$ and $\Phi : (\otimes) \otimes \rightarrow \otimes(\otimes)$ obey respectively hexagon and pentagon identities. In addition there is a trivial object $\underline{1}$ and associated functorial isomorphisms of identity with respect to $\otimes$. A braided group [4][5] is first of all a unital algebra B in such a category, defined in the obvious way except that the identity must be viewed as a morphism $\eta : \underline{1} \rightarrow B$. In addition, we require a coalgebra structure $\Delta : B \rightarrow B \otimes B$ (the coproduct) and $\epsilon : B \rightarrow \underline{1}$ (the counit) obeying axioms which are the same as for an algebra but with arrows reversed. These Δ, ϵ are required to be algebra homomorphisms, where $B \otimes B$ has the braided tensor product algebra structure. Its product $(B \otimes B) \otimes (B \otimes B) \rightarrow B \otimes B$ consists of Ψ to interchange the middle factors followed by products $\cdot$ in B , as well as Φ as needed to order brackets. Finally, there is an antipode or ‘linearized inverse’ morphism $S : B \rightarrow B$ obeying

$$(5) \quad \cdot \circ (\text{id} \otimes S) \circ \Delta = \cdot \circ (S \otimes \text{id}) \circ \Delta = \eta \circ \epsilon.$$

The theory of braided groups can be developed similarly to that of quantum groups or Hopf algebras. A useful (but not necessary) tool is to use a diagrammatic notation as follows. One omits Φ and brackets. One denotes $\otimes$ by juxtaposition, the braiding by $\Psi = \bowtie$ and its inverse by $\Psi^{-1} = \bowtie^{-1}$. Algebraic operations are then drawn as ‘flow charts’ flowing from the top of the page downwards. General morphisms such as S are drawn as nodes. All of this is as for any braided category. Finally, we denote the product and coproduct of B by $\Delta = \wedge$ and $\cdot = \vee$. The principal

axioms of a braided group are shown in this notation in Figure 1. One of the key lemmas that we will need is:

Lemma 2.1. [5] *S is braided-antimultiplicative and anticomultiplicative,*

$$S \circ \cdot = \cdot \circ \Psi \circ (S \otimes S), \quad \Delta \circ S = (S \otimes S) \circ \Psi \circ \Delta.$$

We will also need that if B is a braided group in C with S invertible then

$$(6) \quad \Delta^{op} = \Psi^{-1} \circ \Delta, \quad S^{op} = S^{-1}$$

with the original algebra and counit provide another braided group structure B^{op} in the category C^{op} defined as C but with Ψ, Ψ^{-1} interchanged (i.e. with reversed braid crossings).

In our elementary application we work in the category C_q of $\mathbb{Z}$ -graded vector spaces V, W , etc. with its usual $\otimes$ and (trivial) Φ , but with

$$(7) \quad \Psi_{V,W}(v \otimes w) = w \otimes v \, q^{\deg(v)\deg(w)}$$

extended linearly. Morphisms are degree preserving maps. We work over any field k . The unit object is $\underline{1} = k$.

The algebra of polynomials $B = k[x]$ lives in this category (all its structure maps are morphisms) with $\deg(x) = 1$. We make it into a braided group (the ‘affine braided line’) with

$$\Delta x = x \otimes 1 + 1 \otimes x, \quad \epsilon x = 0, \quad Sx = -x.$$

This looks like the usual additive group Hopf algebra structure on an affine line but Δ extends differently as a homomorphism to the braided tensor product. The braided tensor algebra $B \otimes B$ is generated by $y \equiv x \otimes 1$ and $x \equiv 1 \otimes x$ with the relations $xy = qyx$ (the so-called ‘quantum plane’). Then

$$\Delta f(x) = f(y + x), \quad \epsilon f(x) = f(0),$$

which exhibits Δ as the operation of addition with respect to braided variables where independent x, y q -commute (to emphasise this one may write $f(y +_q x)$, which is the notation indicated in (3-(4)). Working directly from the braided anti/homomorphism properties of S and Δ one also has explicitly

$$(8) \quad \Delta x^i = \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q x^j \otimes x^{i-j}, \quad S(x^i) = q^{\frac{i(i-1)}{2}} (-x)^i,$$

where

$$(9) \quad \begin{bmatrix} i \\ j \end{bmatrix}_q = \frac{(q^i - 1) \cdots (q^{i-j+1} - 1)}{(q^j - 1) \cdots (q - 1)} = \frac{[i]_q!}{[j]_q! [i-j]_q!}.$$

Clearly, the elementary properties of braided groups mentioned above can easily be verified directly from these definitions and boil down to a number of useful identities for q -binomial coefficients, not all of them obvious.

The braided line can be viewed as the ‘correct’ setting for q -analysis in one variable including Jackson’s q -derivative and q -integration[8], q -exponentials, as well as a new theory of q -Fourier transforms[9][10]. Such methods in particular solved the long-standing problem of extending q -differential calculus to the multivariable case on affine q -planes (these turn out to be additive braided groups).

3. INVERTING THE MAP FROM $P[x]$ TO $L(x)$

As in [2] we write (1) as

$$L(x) \equiv \sum_{j=0}^d \frac{L_j}{(q^j - 1) \cdots (q - 1)} x^j = \sum_{j=0}^d \sum_{i=j}^d P_i \frac{(q^i - 1) \cdots (q^{i-j+1} - 1)}{(q^j - 1) \cdots (q - 1)} q^{\frac{j(j-1)}{2}} x^j$$

or, changing the order of summation and using the q-binomial notation,

$$(10) \quad L(x) = \sum_{i=0}^d P_i \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q q^{\frac{j(j-1)}{2}} x^j.$$

Now consider the algebra homomorphism $\phi : k[x] \rightarrow k$ which ‘evaluates at 1’,

$$(11) \quad \phi(x^m) = 1, \quad \forall m.$$

Using this and the structure of the braided line recalled above, we clearly have

$$(12) \quad \begin{aligned} \bar{L}(x) &= \sum_{i=0}^d P_i \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q q^{\frac{j(j-1)}{2}} (-x)^j \\ &= (\phi \otimes S) \sum_{i=0}^d P_i \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q x^{i-j} \otimes x^j = (\phi \otimes S) \circ \Delta P(x) \end{aligned}$$

which is the precise meaning of (3).

Lemma 3.1. *The linear operator $T = (\phi \otimes S) \circ \Delta$ which maps $P(x)$ to $\bar{L}(x)$ has inverse*

$$T^{-1} = (\phi \circ S \otimes \text{id}) \circ \Delta \circ S^{-1}.$$

Proof. The proof is shown in Figure 2 in the diagrammatic notation. The composition $T^{-1} \circ T$ is depicted in the first line in Figure 2(c). Here T^{-1} is linear so we can move the output of ϕ in T to the end of the computation as shown (remember that juxtaposition at the bottom of the line denotes tensor product). We cancel $S^{-1} \circ S$, then use coassociativity of Δ . Finally we use that ϕ is an algebra homomorphism and cancel an ‘antipode loop’ using the axiom of an antipode. The resulting counit combines with Δ to the identity (the counity axiom). Note that ϕ is not a morphism in C_q but we do not use this (we apply ϕ only at the end of each diagrammatic expression). Similarly for the lower line of Figure 2(c). Since these techniques may be unfamiliar, we also write the proof more conventionally. Thus,

$$\begin{aligned} T^{-1} \circ T &= (\phi \circ S \otimes \text{id}) \circ \Delta \circ (\phi \otimes \Delta) = (\phi \otimes \phi \circ S \otimes \text{id}) \circ (\text{id} \otimes \Delta) \circ \Delta \\ &= (\phi \otimes \phi \circ S \otimes \text{id}) \circ (\Delta \otimes \text{id}) \circ \Delta = (\phi \circ \cdot \circ (\text{id} \otimes S) \circ \Delta \otimes \text{id}) \circ \Delta \\ &= \phi(1)(\epsilon \otimes \text{id}) \circ \Delta = \text{id} \end{aligned}$$

and similarly for $T \circ T^{-1}$. These proofs amount to a series of q-identities among q-binomial coefficients by writing out Δ, S explicitly on monomials x^m . $\diamond$

It remains only to compute T^{-1} explicitly. We write $\bar{L} = \sum_{i=0}^d \bar{L}_i x^i$ (without any factorial factors). Then

$$P(x) = T^{-1}(\bar{L}(x)) = (\phi \circ S \otimes \text{id}) \sum_{i=0}^d \bar{L}_i (-1)^i q^{\frac{-i(i-1)}{2}} \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q x^{i-j} \otimes x^j$$




FIGURE 2. (a) The operator T sending P to L and (b) its inverse as proven in (c)

$$= \sum_{i=0}^d \bar{L}_i (-1)^i q^{\frac{-i(i-1)}{2}} \sum_{j=0}^i (-1)^{i-j} q^{\frac{(i-j)(i-j-1)}{2}} \begin{bmatrix} i \\ j \end{bmatrix}_q x^j$$

giving

$$(13) \quad P(x) = \sum_{i=0}^d \bar{L}_i \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q q^{\frac{-j(j-1)}{2}} q^{-j(i-j)} (-x)^j.$$

We can then put in the form of $\bar{L}_i$ in terms of L_i and reorder the summation to extract the P_i . Thus

$$P(x) = \sum_{j=0}^d x^j q^{\frac{-j(j-1)}{2}} \sum_{i=j}^d L_i \frac{(-1)^{i-j} q^{-j(i-j)}}{(q^j - 1) \cdots (q - 1) (q^{i-j} - 1) \cdots (q - 1)}$$

$$= \sum_{j=0}^d x^j \frac{q^{\frac{-j(j-1)}{2}}}{(q^j - 1) \cdots (q - 1)} \sum_{i=j}^d L_i \frac{(-1)^{i-j}}{(q^i - q^j) \cdots (q^{j+1} - q^j)}$$

or P_i as stated in Proposition 1.1. This completes our proof of that.

4. DUALITY UNDER $q \leftrightarrow q^{-1}$

Using one of the basic results from braided group theory, namely the second part of Lemma 2.1, we can also write

$$(14) \quad T^{-1} = (\phi \otimes S^{-1}) \circ \Psi^{-1} \circ \Delta,$$

which is the other expression shown in Figure 2(b) and the precise meaning of (4). This computes more directly as (13), with $q^{-j(i-j)}$ coming from the braiding $\Psi^{-1}(x^j \otimes x^{i-j})$ and $q^{\frac{-j(j-1)}{2}}$ from S^{-1} .

Corollary 4.1. *The operator T^{-1} is exactly the same as the operator T but with q replaced by q^{-1} .*

Proof. We see that T^{-1} has just the same form as T but with Δ^{op} and S^{op} in place of Δ, S . This is the opposite braided group structure on $k[x]$, namely the same form on degree 1 but developed as a braided group in the category $C_q^{\text{op}} = C_{q^{-1}}$. One may also see this explicitly by observing that

$$\begin{bmatrix} i \\ j \end{bmatrix}_{q^{-1}} = q^{-j(i-j)} \begin{bmatrix} i \\ j \end{bmatrix}_q$$

so that (13) can be written as

$$(15) \quad P(x) = \sum_{i=0}^d \bar{L}_i \sum_{j=0}^i (-1)^j \begin{bmatrix} i \\ j \end{bmatrix}_{q^{-1}} q^{\frac{-j(j-1)}{2}} (-x)^j$$

which has the same form as our original (12). $\diamond$

Finally, we mention another way to obtain the above duality using braided group theory, this time in the form of ‘q-analysis’. Namely, the displacement of a function by 1 can be written as a q-exponential of q-derivatives (the braided Taylor’s theorem[9]). In the present one-variable case it is easily verified on monomials and takes the form

$$(\phi \otimes \text{id})\Delta = e_q^{\partial_q} = \sum_{i=0}^{\infty} \frac{(\partial_q)^i}{[i]_q!}$$

acting on polynomials (the sum is only finite). Here

$$\partial_q f(x) = \frac{f(qx) - f(x)}{x(q-1)}$$

is Jackson’s q -derivative[8]. (Also in our case one can cancel the $(q-1)$ factors here and in $[i]_q!$ if one prefers.) Moreover,

$$e_q^A e_{q^{-1}}^{-A} = 1$$

for any nilpotent operator A as one may see by q-differentiating[6, Lem. 3.2.2]. Hence

$$T = S \circ e_q^{\partial_q} = e_q^{-\partial_{q^{-1}}} \circ S.$$

The second equality here is easily verified on monomials. Then

$$T^{-1} = e_{q^{-1}}^{-\partial_q} \circ S^{-1} = S^{-1} \circ e_{q^{-1}}^{\partial_{q^{-1}}},$$

which has the same form with q and q^{-1} interchanged.

REFERENCES

- [1] N. Boston, W. Dabrowski, T. Foguel, P.J. Gies, J. Leavitt, D.T. Ose and D.A. Jackson. The proportion of fixed-point-free elements of a transitive permutation group. *Commun. Algebra* 21:3259–3275, 1993.
- [2] P.J. Cameron. Seminar, November 2000 (unpublished).
- [3] R.P. Stanley. *Enumerative Combinatorics, Vol 1*. Cambridge University Press, 1997.
- [4] S. Majid. Braided groups. *J. Pure and Applied Algebra*, 86:187–221, 1993.
- [5] S. Majid. Algebras and Hopf algebras in braided categories. volume 158 of *Lec. Notes in Pure and Appl. Math*, pages 55–105. Marcel Dekker, 1994.
- [6] S. Majid. *Foundations of Quantum Group Theory*. Cambridge Univeristy Press, 1995.
- [7] A. Joyal and R. Street. Braided monoidal categories. Mathematics Reports 86008, Macquarie University, 1986.
- [8] F.H. Jackson. q -Integration. *Proc. Durham Phil. Soc.*, 7:182–189, 1927.
- [9] S. Majid. Free braided differential calculus, braided binomial theorem and the braided exponential map. *J. Math. Phys.*, 34:4843–4856, 1993.
- [10] A. Kempf and S. Majid. Algebraic q -integration and Fourier theory on quantum and braided spaces. *J. Math. Phys.*, 35:6802–6837, 1994.

SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY, UNIVERSITY OF LONDON, 327 MILE END RD, LONDON E1 4NS, UK